

To the Editor—T Chan and colleagues should be congratulated on their report of the stroke-like episode following a rapid ascent to altitude.¹ There are few points concerning differential diagnosis that are worth considering. Firstly, stroke or stroke-like episodes have been described in association with high altitude since 1895.² Thromboembolic aetiology is a possibility, although absence of risk factors and stay shorter than reported by Jha et al³ may argue against that. The authors might also consider the possibility of a cryptogenic stroke due to paradoxical embolism. Consequent to hypoxic pulmonary vasoconstriction encountered at altitude, there is a rise in pulmonary vascular resistance and arterial pressure and increased likelihood of right to left shunting. Depending on the population studied, some 20 to 30% of individuals have an asymptomatic patent foramen ovale.⁴ This, in the presence of a deep vein thrombosis, could provide a substrate for stroke. Secondly headache is not a usual feature of an ischaemic event. It is, however, a symptom of acute mountain sickness or high-altitude cerebral oedema and can also occur in posterior reversible encephalopathy syndrome (PRES) or migraine.

While PRES is characterised by vasogenic oedema, it can precipitate intense vasospasm and cerebral infarction. Admittedly the patient has had no classical risk factors for PRES, but the distribution of lesions hints at this possibility.⁵ Transient focal neurology occurring at altitude in patients with a history of migraines has also been reported.^{6,7} Absence of radiologically evident vascular pathology and a degree of reversibility observed could reflect a vasospastic aetiology.

Piotr Szawarski, FRCA, MSc

Email: piotr.szawarski@gmail.com

Department of Anaesthesia, Guy's & St Thomas' NHS Foundation Trust, Great Maze Pond, London SE1 9RT, United Kingdom

Emily W Y Tam, BM, FRCR

Department of Radiology, Medway Maritime Hospital, Windmill Road, Gillingham ME7 5NY, United Kingdom

Paul Richards, MRCCGP, MSc

The Surgery and Travel Clinic, London Road, Wickford, Essex, United Kingdom

References

1. Chan T, Wong WW, Chan JK, Ma JK, Mak HK. Acute ischaemic stroke during short-term travel to high altitude. *Hong Kong Med J* 2012;18:63-5.
2. West JB, Schoene RB, Milledge JS. High altitude medicine and physiology. 4th ed. London: Hodder Arnold; 2007: 312.
3. Jha SK, Anand AC, Sharma V, Kumar N, Adya CM. Stroke at high altitude: Indian experience. *High Alt Med Biol* 2002;3:21-7.
4. Wu LA, Malouf JF, Dearani JA, et al. Patent foramen ovale in cryptogenic stroke: current understanding and management options. *Arch Intern Med* 2004;164:950-6.
5. Bartynski WS. Posterior reversible encephalopathy syndrome, part 1: fundamental imaging and clinical features. *AJNR Am J Neuroradiol* 2008;29:1036-42.
6. Jenzer G, Bärtsch P. Migraine with aura at high altitude. *J Wilderness Med* 1993;4:412-5.
7. Murdoch DR. Focal neurological deficits and migraine at high altitude. *J Neurol Neurosurg Psychiatry* 1995;58:637.

Rational emergency stroke care in Hong Kong

To the Editor—Dr Eric Chan¹ makes the excellent point that we need to concentrate interventional stroke services in a small number of well-chosen locations within Hong Kong. This is particularly so given its relative rarity and the expertise required to achieve consistent high standards in this difficult therapeutic area. We have previously suggested that speed is of the essence in the delivery of acute interventional stroke care in Hong Kong.²

As he states, Hong Kong has an excellent road network, which means that both primary diversion of critically ill patients and interfacility transfers are very feasible and have been shown to be safe.^{3,4}

As previously suggested by us, these principles should also be extended to the management of acute ST elevation myocardial infarction in Hong Kong.⁵ Our professional ambulance personnel in Hong Kong would be perfectly capable of triaging appropriate