

# Reversal of warfarin in life-threatening bleeding

Hong Kong Med J 2015;21:191

DOI: 10.12809/hkmj154509

*To the Editor*—I read with interest the article “The benefit of prothrombin complex concentrate in decreasing neurological deterioration in patients with warfarin-associated intracerebral haemorrhage” by Fong et al in the December 2014 issue of the *Hong Kong Medical Journal*.<sup>1</sup> The Australian guidelines for warfarin reversal quoted in the article were outdated.<sup>2</sup> The most recent Australian guidelines can be found in the March 2013 issue of the *Medical Journal of Australia*.<sup>3</sup>

**Daniel SK Ng \***, FACEM, FHKAM (Emergency Medicine)

Emergency Department, Swan District Hospital, 1 Eveline Road, Middle Swan, Western Australia 6056

\* Corresponding author: daniel.ng@health.wa.gov.au

## References

1. Fong WC, Lo WT, Ng YW, et al. The benefit of prothrombin complex concentrate in decreasing neurological deterioration in patients with warfarin-associated intracerebral haemorrhage. *Hong Kong Med J* 2014;20:486-94.
2. Baker RI, Coughlin PB, Gallus AS, Harper PL, Salem HH, Wood EM; Warfarin Reversal Consensus Group. Warfarin reversal: consensus guidelines, on behalf of the Australasian Society of Thrombosis and Haemostasis. *Med J Aust* 2004;181:492-7.
3. Tran HA, Chunilal SD, Harper PL, Tran H, Wood EM, Gallus AS; Australasian Society of Thrombosis and Haemostasis (ASTH). An update of consensus guidelines for warfarin reversal. *Med J Aust* 2013;198:198-9.