

To the Editor—We read with interest the case report describing the ‘oesophageal haemangioma’.¹ This related to a 59-year-old woman with a vascular lesion in her oesophagus. The discussion refers to the lesion as a vascular tumour. The question that we would like to raise is whether this was a vascular tumour or a vascular malformation. New generations of medical students are being specifically taught to recognise and appreciate the difference. We no longer want to find young adults with disfiguring facial lesions waiting for their ‘haemangiomas’ to involute spontaneously.

The current teaching in both medical schools is based on the classification system adopted by the International Society for the Study of Vascular Anomalies (ISSVA).² This basic classification was adopted by the ISSVA in 1996 and distinguishes between the two main types of vascular anomalies, namely the vascular tumour and the vascular malformation. Vascular tumours grow by cellular hyperplasia, most usually involving endothelial cells. Vascular malformations, however, have a quiescent endothelium and are regarded as localised anomalies in vascular anatomy, most probably due to faulty

development in utero.

The use of a consistent classification system across the disciplines would be helpful and this particularly includes the radiological and pathological descriptions of vascular lesions. Proper classification will also guide us in applying the appropriate therapy. Medicine evolves and terminology becomes more precise. Is it possible that ‘oesophageal haemangiomas’ do not really exist? Perhaps not in the world of Mulliken and Glowacki,³ and most probably not according to the ISSVA. Is it time to put them to rest in Hong Kong?

Andrew Burd, MD, FRCS

Email: andrewburd@surgery.cuhk.edu.hk

Division of Plastic, Reconstructive and Aesthetic Surgery, Department of Surgery, The Chinese University of Hong Kong, Prince of Wales Hospital, Hong Kong

Godfrey CF Chan, DMD, MD

Division of Paediatric Haematology/Oncology, Department of Paediatrics and Adolescent Medicine, The University of Hong Kong, Queen Mary Hospital, Hong Kong

References

1. Rasalkar DD, Chiu PW, Teoh AY, Chu WC. Oesophageal haemangioma: imaging characteristics of this rare condition. *Hong Kong Med J* 2010;16:230-1.
2. Introduction: ISSVA classification. In: Color atlas of vascular tumors and vascular malformations. Enjolras O, Wassef M, Chapot R, editors. New York: Cambridge University Press; 2007: 1-11.
3. Mulliken JB, Glowacki J. Hemangiomas and vascular malformations in infants and children: a classification based on endothelial characteristics. *Plast Reconstr Surg* 1982;69:412-22.