

## References

1. Ch'in KY, Tang MY. Congenital adenomatoid malformation of one lobe of a lung with general anasarca. Arch Pathol 1949;48:221-9.
2. Stocker JT, Madewell JE, Drake RM. Congenital cystic adenomatoid malformation of the lung. Hum Pathol 1977;8:156-71.
3. Moerman P, Fryns JP, Vandenberghe K, Devlieger H, Lauweryns JM. Pathogenesis of congenital cystic adenomatoid malformation of the lung. Histopathology 1992 Oct;21(4):315-21.
4. Miller RK, Sieber WK, Yunis EJ. Congenital adenomatoid malformation of the lung: a report of 17 cases and review of the literature. Pathol Annu 1980;15:378-402.
5. Neilson IR, Russo P, Laberge JM, et al. Congenital adenomatoid malformation of the lung: current management and prognosis. J Pediatr Surg 1991 Aug;26(8):975-80.
6. Han YM, Lee DK, Lee SY, et al. Adult presentation of congenital cystic adenomatoid malformation of the lung: a case report. J Korean Med Sci 1994 Feb;9(1):86-91.
7. Patz EF, Jr, Muller NL, Swensen SJ, Dodd LG. Congenital cystic adenomatoid malformation in adults: CT findings. J Comput Assisted Tomogr 1995 May/Jun;19(3):361-4.
8. Murphy JJ, Blair GK, Fraser GC, et al. Rhabdomyosarcoma arising within congenital pulmonary cysts: report of three cases. J Pediatr Surg 1992 Oct;27(10):1364-7.
9. Benjamin DR, Cahill JL. Bronchioalveolar carcinoma of the lung and congenital cystic adenomatoid malformation. Am J Clin Pathol 1991 Jun;95(6):889-92.
10. Harrison MR, Adzick NS, Jennings RW. Antenatal intervention for congenital cystic adenomatoid malformation. Lancet 1990;336:965-7.

HKMJ 1996;2:201-206

## PICTORIAL MEDICINE



This is the retinal appearance of a 20-year-old man who presented with a total white cell count of  $341 \times 10^9/L$  and a platelet count of  $29 \times 10^9/L$ . He was subsequently diagnosed as having acute biphenotypic leukaemia. He complained of reduced visual acuity and fundoscopic examination showed multiple retinal haemorrhages and Roth's spot. The latter are thought to be due to leucostasis causing central ischaemia with surrounding haemorrhages.

*Submitted by W Yeo, N Wickham, PJ Johnson.*

*Department of Clinical Oncology, the Chinese University of Hong Kong,*

*Prince of Wales Hospital, Shatin, Hong Kong.*