



Supplementary material

The supplementary material was provided by the authors and some information may not have been peer reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by the Hong Kong Academy of Medicine and the Hong Kong Medical Association. The Hong Kong Academy of Medicine and the Hong Kong Medical Association disclaim all liability and responsibility arising from any reliance placed on the content.

Supplement to: CHM Wong, ICH Ko, DKW Leung, et al. Anaemia and iron deficiency in advanced prostate cancer: revisiting a common morbidity. Hong Kong Med J 2025 Apr;31(2):154-8 | Epub 8 Apr 2025. <https://doi.org/10.12809/hkmj2411653>.

SUPPLEMENTARY TABLE 1. Multivariate analysis of factors contributing to anaemia in a cohort of prostate cancer patients

	Not castrated	Castrated	P value
Likelihood of anaemia	0.3421	0.6818	0.002
Odds ratio	1.99		
	Non-metastatic	Metastatic disease	P value
Likelihood of anaemia	0.4	0.7188	0.004
Odds ratio	1.78		
	Non-CRPC	CRPC	P value
Likelihood of anaemia	0.4671	0.5035	0.151
Odds ratio	1.08		
	Normal albumin	Hypoalbuminaemia	P value
Likelihood of anaemia	0.4848	1.0	0.015
Odds ratio	2.05		
	No CKD	CKD	P value
Likelihood of anaemia	0.4643	0.6538	0.112
Odds ratio	1.41		

Abbreviations: CKD = chronic kidney disease; CRPC = castration-resistant prostate cancer

SUPPLEMENTARY TABLE 2. Associations between haemoglobin level and testosterone level

	Effect coefficient	P value
Overall	-0.025	0.825
Subgroup		
Localised disease	0.323	0.024
Metastatic hormone-sensitive disease	-0.081	0.734
Castration-resistant status	0.365	0.269